


### Clinical severity of COVID cases diagnosed at UHS

2021-12-26 to 2022-04-16



Adapted from the NIH COVID-19 Treatment Guidelines:

- I. **Asymptomatic** - Individuals who test positive for SARS-CoV-2 using a virologic test (i.e., a nucleic acid amplification test [NAAT] or an antigen test) but who have no symptoms that are consistent with COVID-19.
- II. **Mild** – Individuals who have any of the various signs and symptoms of COVID-19 (e.g., fever, cough, sore throat, malaise, headache, muscle pain, nausea, vomiting, diarrhea, loss of taste and smell) without shortness of breath, dyspnea, or abnormal chest imaging.
- III. **Moderate** – Individuals who show evidence of lower respiratory disease during clinical assessment or imaging (e.g., shortness of breath, dyspnea, or abnormal chest imaging) and who have an oxygen saturation (SpO<sub>2</sub>) ≥94% on room air at sea level.
- IV. **Severe** – Individuals who have SpO<sub>2</sub> <94% on room air at sea level, a ratio of arterial partial pressure of oxygen to fraction of inspired oxygen (PaO<sub>2</sub>/FiO<sub>2</sub>) <300 mm Hg, a respiratory rate >30 breaths/min, or lung infiltrates >50%.
- V. **Critical** – Individuals who have respiratory failure, septic shock, and/or multiple organ dysfunction.

1 Source: <https://www.covid19treatmentguidelines.nih.gov/overview/clinical-spectrum/>

		I		II		III		IV		V		?		Total
Week	Dates	Count	I%	Count	II%	Count	III%	Count	IV%	Count	V%	Count	?%	Count
16	4/10/22 - 4/16/22	14	15.4%	71	78.0%	3	3.3%	0	0%	0	0%	3	3.3%	91